

Patient Name: _____

Date: _____

Circle appropriate score for each section and total the score below.

CRITERIA	SCORE	
	Value	Patient's Score

History (2, 4, 8, 11, 12, 13)

History of Falls in past 12 months	Self-reported number of falls or near falls in the past 12 months	None	1	
		1	2	
		>2	3	
Medications taken in past 7 days	Medications include PCA/opioids, benzodiazepines, anticonvulsants, anti-hypertensives, diuretics, hypnotics, laxatives, sedatives and psychotropics	None	1	
		1 to 2	2	
		>3	3	
History of Lateral Ankle Sprains in past 12 months	Self-reported number of LAS in the past 12 months	None	1	
		> 1	3	

Physical (3, 6, 7, 10, 12)

Pronation	In static standing, does the patient present with greater than 5 degrees of calcaneal valgus, collapsed MLA and/or navicular drop	WNL	1	
		Excessive pronation	2	
Range of Motion	Range of motion in dorsiflexion, hip extension, internal rotation and adduction.	WNL	1	
		Limited ROM	2	
Strength / Endurance	Chair-Stand Test: Have patient sit in chair, place hands on opposite shoulders, keep feet flat on floor, keep back straight with arms against chest. On "Go", rise to full standing position, then sit back down. Repeat for 30 seconds. Record number of times patient stands. *See Scoring Below*	WNL	1	
		Decreased strength / Endurance	2	

Proprioception (1, 5, 9, 10, 12)

Vision	Ability to see objects clearly, judging distances, difficulty seeing in reduced light, diagnosed with eye disease interfering with vision	WNL	1	
		Decreased vision	2	
Standing Balance / Vestibular	Standing unsupported with feet touching for 1 minute	> 1 minute	1	
		< 1 minute	2	
		Cannot perform	3	
Standing Balance / Vestibular	Tandom Stand for 10 seconds (One foot in front of the other with heel touching opposite toe)	> 10 seconds	1	
		< 10 seconds	2	
		Cannot perform	3	
Neuropathy	Sensation in the feet or legs (numbness / pins and needles)	WNL	1	
		Decreased sensation	3	

CRITERIA		SCORE		
		Value	Patient's Score	
Function/Gait (10, 11, 12)				
Community Ambulation	Self-reported or observed ability to ambulate (with or without aids) in the community	WNL	1	
		Decreased ability to safely ambulate in community	3	
Transitioning	Self-reported or observed ability to ambulate and transition in the home without holding onto doors or walls	None	1	
		Decreased ability to transition between surfaces/navigate corners and obstacles	3	
Total				

GREEN/LOW RISK (TOTAL: 12-14)	Patient is a low risk for falls. Continue to monitor, minimize risk factors.
YELLOW/MOD RISK (TOTAL: 15 - 20)	Patient is at a moderate risk for falls. Refer for physical therapy and assistive devices such as Surestep Stabilizer.
RED/HIGH RISK (TOTAL: 21-32)	Patient is at a high risk for falls. Refer for fall prevention program, physical therapy and assistive devices such as Surestep Stabilizer.

Chair-Stand Below Average Scores		
Age	Men	Women
60-64	<14	<12
65-69	<12	<11
70-74	<12	<10
75-79	<11	<10
80-84	<10	<9
85-89	<8	<8
90-94	<7	<4

Physician Signature _____ **Date** _____

REFERENCES

- Berg et al. (1992) Balance during standing, sitting to standing, one foot. Can J Pub Health. 83(2):S7-11.
- Boulgarides LK, et al. (2003) Use of clinical and impairment-based tests to predict falls by community-dwelling older adults. Phys Ther. 83(4):328-39.
- Centers for Disease Control and Prevention. (2017) National Center for Injury Prevention and Control. 30-Second Chair Stand. www.cdc.gov/steady.
- Centers for Disease Control and Prevention. (2017) National Center for Injury Prevention and Control. Medications Linked to Falls. www.cdc.gov/steady.
- Centers for Disease Control and Prevention. (2017) National Center for Injury Prevention and Control. The 4-Stage Balance Test. www.cdc.gov/steady.
- Chiacchiero M, et al. (2010) The Relationship Between Range of Movement, Flexibility, and Balance in the Elderly. Topics in Geriatric Rehabilitation. 26(2):147-154.
- Hee Cho K, et al. (2012) Effect of Lower Limb Strength on Falls and Balance of the Elderly. Ann Rehabil Med. 36:386-393.
- Hendrich A. Fall Risk Assessment for Older Adults: the Hendrich II Fall Risk Model. Try This. Best Practices in Nursing Care to Older Adults. Revised 2016 (8).
- Melzer I, et al. (2010) A retrospective analysis of balance control parameters in elderly fallers and non-fallers. Clinical Biomechanics. 25:984-988.
- Phelan EA, et al. (2015) Assessment and Management of Fall Risk in Primary Care Settings. Med Clin North Am. 99(2):281-293.
- Poe SS, et al. (2007) The Johns Hopkins Fall Risk Assessment Tool: postimplementation evaluation. J Nurs Care Qual. 22(4):293-298.
- Russell MA, et al. (2009) Development of the Falls Risk for Older People in the Community (FROP-COM) screening tool. Age Ageing. 38(1):40-46.
- Terada M, et al. (2018) Altered postural control variability in older-aged individuals with a history of lateral ankle sprain. Gait & Posture. 60:88-92.